

Gözdenur ÇAKAR^{1a}

Işıl SARAÇ SİVRİKAYA^{2a}

Ersin KARAKAYA^{3a}

Abdullah GÜLLER^{2b}

¹Bingöl University, Institute of Sciences, Bee and Bee product, Bingöl

²Bingöl University, Faculty of Agriculture, Plant Protection, Bingöl

³Bingöl University, Faculty of Agriculture, Agricultural Economics, Bingöl

^{1a}ORCID: 0000-0002-2359-1312

^{2a}ORCID: 0000-0002-5991-2173

^{3a}ORCID: 0000-0002-6734-4962

^{2b}ORCID: 0000-0003-3887-4208

*Corresponding author:

aguller@bingol.edu.tr

DOI

<https://doi.org/10.46291/ISPECJASv015iss4pp795-805>

Alınış (Received): 25/05/2021

Kabul Tarihi (Accepted): 26/06/2021

Keywords

Lavender essential oil, summer savory essential oil, inhibition, *F. solani*

Evaluation of the *In Vitro* Fungicidal Activity of Summer Savory and Lavender Essential Oils Against *Fusarium solani*

Abstract

Fusarium spp is one of the major phytopathogenic microfungus strains causing severe losses in many economically cultivated crops. The soil-borne pathogen *Fusarium solani* has historically been considered a serious agent across the globe, causing vascular wilt and root rot in agro-economic crops and eventually leading to plant death. Three different concentrations (1 µl, 2 µl, and 4 µl) of essential oils (EO) extracted from lavender (*Lavandula officinalis* L.) and summer savory (*Satureja hortensis* L.) plants were mixed separately with PDA medium, and their antifungal effect against *F. solani* was investigated *in vitro*. When the results of the experiment were evaluated statistically, it was determined that the increasing concentrations of summer savory essential oil repressed the mycelial growth of the fungus, while lavender oil did not have any positive or negative effects. The inhibition activity of summer savory EO on *F. solani* was calculated as 43, 53, and 90% at the concentrations of 1, 2, and 4 µl, respectively. In this study, it was found that summer savory EO, even at a minimum dose, had a negative effect on agriculturally important wilt agent. In this context, it can be asserted that summer savory EO is a promising natural substance for the development of various fungicide solutions to prevent fungal diseases caused by vascular origin.

1. INTRODUCTION

Fusarium solani, a well-known opportunistic agent that can be found in every part of the world with varying subspecies in saprophytic and parasitic forms, is a cause of fungal infection in animals and humans (Brayford, 1993; O'Donnell et al., 2015; Tuxbury et al., 2014). Surviving in the soil for years, the agent can be spread over long distances through wind, agricultural equipment, and irrigation. Host plants infected with different strains of fungus often exhibit irreversible severe symptoms such as wilting, die back, root and fruit rot (Al-Sadi et al., 2014; Bueno et al., 2014; Yaseen and D'Onghia, 2012). With its wide host range on a global scale, *F. solani* has been reported in many countries including Turkey, Czech Republic, Japan, India, Iran, the United States, Iraq, Malaysia, and Germany (Kurt et al., 2020; Ondřej et al., 2008; Sugiura et al., 2003; Ramteke et al., 2019; Abedi-Tizaki et al., 2016; Romberg and Davis, 2007; Hafizi et al., 2013; Farahani-Kofoet et al., 2020). Essential oils, also known as volatile oils, are natural oily substances that are obtained from the leaves, fruits, barks or roots of aromatic plants. They are liquid at room temperature, easily crystallized, usually colorless or light yellow and have a strong odor. Distillation, extraction, and pressing are commonly used methods to obtain these oils. Until today, more than 2000 chemical components have been reported to be present in EOs, mainly terpenes, phenylpropanes and compounds containing small amounts of alcohols, aldehydes, esters, phenols, nitrogen, and sulfur (Çelik and Çelik, 2007; Kılıç, 2008). Due to the physiological effects of the active compounds they have at the cellular level, EOs are widely used singly or in a mixed manner in the fields of aromatherapy and industry. One of the most researched aspects of EOs is related to their antimicrobial and insect-repellent activities. The effectiveness of these oils varies widely depending on the type and amount of the active ingredients they contain (Ceylan,

1983; 1987; Linskens and Jackson, 1997; Bayaz, 2014; Limoncu et al., 2017). Previous studies in the literature provide a wide-ranging documentation on the antimicrobial functionality of EOs and their contents. The antagonistic effect of EOs against many viral, bacterial, and fungal pathogens of plant and animal origin has been well-investigated. The antibacterial effect of EOs on many microorganisms, including Gram (-) and Gram (+) bacteria, has been extensively covered in the literature (Nostro et al., 2000; El-Shazly et al., 2002; Al-Howiriny, 2003; Sartoratto et al., 2004; Chouhan et al., 2017; Gadisa et al., 2019; Man et al., 2019; György et al., 2020). Some other studies have focused on the antiviral activities of EOs. Bammi et al. (1997) and Bishop (1995) reported that EOs were effective inhibitors against the replication of Epstein-Barr virus (EBV) and *Tobacco mosaic virus* (TMV), respectively. Today, there is a comprehensive documentation on the inhibitory effect of EOs against SARS-CoV-2, Influenza A/WS/33, and herpes simplex virus type 1 (HSV-1) infections. The antifungal properties of EOs vary depending on the source and part of the plant, growing environment, climate, and quantity. Many plants in a broad spectrum, including sage, laurel, cinnamon, thyme, lemongrass, nettle, basil and geranium, have natural fungistatic and/or fungicidally effective essential oil compounds that tend to reduce the growth of various fungal pathogens (Angioni et al., 2006; Pinto et al., 2007; Guynot et al., 2003; Wang et al., 2009; Rasooli et al., 2006; Tzortzakis et al., 2007; Sahin et al., 2004; Nguefack et al., 2009). Increasing resistance to fungicides in plants has led to the need for the development of new antifungal strategies. Thanks to their content of terpenoids, terpenes, and aromatic compounds, EOs are one of the main candidates for the development of antifungal preparations (Bassole and Juliani, 2012). In this study, we investigated the anti-fungal activities of lavender and summer savory essential oils against *F.*

solani, one of the main fungal agents that infect economically important crop plants, in increasing doses under laboratory conditions.

2. MATERIALS and METHODS

2.1. Experimental fungal isolate and source of essential oils

From previous studies, a fungus (*F. solani*) isolated from the diseased parts of bean plants grown in Bingol province was used as the experimental pathogen (Sarac Sivrikaya et al. 2021). The essential oils of lavender and summer savory plants used in the experiments were extracted by a commercial enterprise through the water vapor distillation method using the Clevenger apparatus. The essential oils were kept in dark-colored and tightly closed bottles at + 4 °C until use.

2.2. Culture media for inhibition tests

A common medium Potato Dextrose Agar (PDA) was used for the purification and growth of the fungal pathogen. The prepared medium was sterilized, and approximately 20 ml of liquid medium was transferred to glass petri dishes. Immediately, 1 µl, 2 µl, and 4 µl of essential oils were added to the PDA media to form application groups. Each application was carried out in 3 repetitions, as well as the control groups with only the pathogen and PDA medium without essential oil.

2.3. Preparation of pathogen fungus for inoculum

F. solani was inoculated into the prepared media and incubated for 7 days at 25 °C for growth. After fungal growth, 8 mm diameter fungal discs were cut out from the culture medium to test the antifungal activity of the oils.

2.4. Assessment of antifungal activity and statistical analysis

To determine fungal inhibition, the removed fungal discs were placed in the center of the petri dishes containing PDA medium, and the petri dishes were covered with parafilm to prevent a possible contamination. After 7 days of incubation at 24 ± 1°C, the diameters of fungal colonies

were measured vertically and in separate directions ignoring 8 mm fungal disc diameters, photographed and recorded (Benjilali *et al.*, 1984). Percent inhibition was calculated according to the formula reported Deans and Svoboda (1990).

$$\text{Inhibition rates (\%)} = \frac{gc-gt}{gc} \times 100$$

Analysis of variance (ANOVA) was used to determine the differences between the treatment groups. It was tested whether the variances of the data were homogeneous, and the means were compared using the DUNCAN test. The statistical significance was set at P<0.05.

3. RESULTS and DISCUSSION

F. solani is one of the most common fungus species in agricultural production areas worldwide (Saremi et al., 2011). Soil-borne pathogen attack causes plant death due to the development of root rot complex on field and horticultural crops, resulting in serious yield losses. Recently, it has been reported by most researchers that essential oils obtained from various medicinal and aromatic plants can be used as natural fungicides in the control of some plant and bee diseases (Djihane et al., 2017; Tutun et al., 2018). Many plant species can synthesize species-specific EOs in different amounts at different developmental stages in different tissues. The biological activities of essential oils vary considerably depending on the genotype and ecological environment of the plant and the test method and microorganism used (Rota et al., 2004; Yeşil Çelikleş et al., 2007). Thanks to their chemical composition, essential oils obtained from plants such as sage, rosemary, laurel, oregano, clove, thyme, cumin, lavender, lemongrass, marjoram, mint, lemon balm, eucalyptus, fennel, and cinnamon have been previously used to control plant fungal diseases (Soylu et al., 2005; Lee et al., 2007). In this study, we investigated the antifungal effects of lavender and summer savory EOs containing chemical aromatic compounds with antimicrobial properties. Under *in vitro* conditions, the inhibitory effect of

three different doses (1 μ l, 2 μ l, 4 μ l) was examined against mycelial growth of *F. solani*, and their percentage inhibition values were determined (Table 1). Summer savory EOs clearly suppressed the mycelial growth of the test pathogen *F. solani*, but

lavender EOs did not generate any noticeable results. Summer savory oil showed a biological activity at increasing doses, which was confirmed both statistically and by the radial inhibition zone (Figure 1).

Table 1. *In vitro* antifungal activity of *S. hortensis* and *L. officinalis* essential oils against *F. solani*

Extracts types	Dose	Inhibition (%)	Mean growth diameter (mm)
<i>S. hortensis</i>	4 μ l	90.0	10,75 ^{a*}
	2 μ l	53.0	33,25 ^b
	1 μ l	43.0	39,00 ^c
<i>L. officinalis</i>	4 μ l	3.0	60,75 ^d
	2 μ l	3.0	61,00 ^d
	1 μ l	3.0	62,00 ^{de}
NC**	-	0.0	62,75 ^e

*The difference between the means with the same letters in the same column is not statistically significant ($p < 0.05$) **NC: no-treatment control



Figure 1. The antifungal activities of 4 μ l of *S. hortensis* and *L. officinalis* EOs against *F. solani* isolated from *P. vulgaris* in glass petri dishes containing PDA medium. A: *S. hortensis* EOs treatment, B: *L. officinalis* EOs treatment; C: NC (No-treatment control)

Lavender, belonging to the Lamiaceae family, has two main important antimicrobial chemical compounds: linalyl acetate and linalool (Śmigielski et al., 2013). Due to their antibacterial, antifungal, and healing properties, EOs extracted from some species of genus *Lavandula* are one of today's popular topics. Lavender EOs have been extensively used as a strong antimicrobial agent to eliminate foodborne pathogens, such as *Escherichia coli*, *Salmonella typhimurium*, *Staphylococcus aureus*, *S. epidermidis*, *Bacillus cereus*, and *Listeria monocytogenes*, and *L. innocua* as

well as other bacteria (*Clostridium perfringens*, *Pseudomonas aeruginosa*, *Proteus vulgaris*, *Enterobacter aerogenes*) (Dadalioglu and Evrendilek, 2004; Nikolis et al., 2014; Djenane et al., 2012; Varona et al., 2013; Viuda-Martos et al., 2011). Furthermore, some previous studies on lavender EO focused on its inhibitory effect against human fungal strains including *Candida albicans* and *Aspergillus niger*, and while some others focused on controlling various phytopathogenic fungi (Tarek et al., 2014). Yohalem and Passey, (2011) reported that *L. angustifolia* and *L. x*

intermedia EOs were toxic to the wilting agent *Verticillium dahlia* in the strawberry plant. The essential oil from *L. stoechas*, another species of the genus Lavender, was reported to have an effect as a biofungicide by negatively affecting the membrane integrity of the gray mold (*Botrytis cinerea*) and wilting pathogen (*V. dahlia*) in tomato plants (Kadoglidou et al., 2011; Soylu et al., 2010). Additionally, a different study showed that EO from *L. stoechas* was effective against *Phytophthora infestans*, which is responsible for late blight disease in tomato (Soylu et al., 2006). In contrast, other reports showed that EO extracted from *L. angustifolia* had a moderate antifungal effect against *B. cinerea*, *Rhizopus stolonifer*, and *A. niger*, which are critical fungal pathogens in the postharvest stage of strawberries (Hadian et al., 2008). Similarly, Lopez-Reyes et al. (2010) showed that essential oil from *L. officinalis* exhibited a low toxicity against *B. cinerea* and *Penicillium expansum* fungal infection in postharvest apples, which is consistent with the results obtained for *L. officinalis* EO against *F. solani* in the present study. As can be seen in the abovementioned studies, lavender essential oils were reported to have a unique fungicidal activity against different fungi species depend on the lavender species from which they were extracted. But, the results of our study showed that lavender oil did not exhibit any statistically significant toxicity against soil-borne fungus *F. solani*, even in the increasing doses. In a previous study, the low antifungal efficacy of *L. hybrida* and *L. angustifolia* essential oils was attributed to the insufficiency of thymol and carvacrol components (Giordani et al., 2004). Possibly, the same applies to our study, and therefore, antifungal activity could not be determined at the doses used in the present study. A striking point to note in Figure 1 is that there is a minimal difference between the inhibition effects of the lavender essential oil applications. In this context, it is necessary to apply a higher concentration than we used in order to observe the

inhibition zone against *F. solani* fungus. Al-Naser and Al-Abrass (2014) support this idea in that they used the same essential oil, but obtained by the GC-MC method, and found a broad spectrum fungistatic activity against *F. solani*, *F. oxysporum*, and *A. niger*, possibly due to the fact that they used higher amounts of essential oil (500, 1000, and 1500 μ l) than ours (1, 2, and 4 μ l). As a result, most reports have supported that lavender EOs can be used, to some extent, as an alternative to the synthetic fungicides being used currently. The summer savory-associated antifungal outputs were reported to be more remarkable than the others, and several previous studies supported the potential antifungal activity of *Satureja hortensis*. We found that the test pathogen was more susceptible to summer savory EOs compared to lavender oil, even in minimal doses. Summer savory essential oil contains active ingredients with antimicrobial and food preservative properties, mainly phenol derivative carvacrol, cyclohexone, cymene, and thymol (Özkalp and Özcan, 2009). Şahin et al. (2003) tested hexane and methanol extracts of summer savory against 55 bacterial species, and 31 isolates of 1 yeast and 4 fungus species and reported that Hexane extracts were toxic to a wide range of test microorganisms, but both extracts did not display inhibition against phytopathogenic fungi *Alternaria alternata*, *A. flavus*, *Fusarium oxysporum*, and *Penicillium* spp, suggesting that EOs have more fungicidal activity than extracts. Like our study, Dikbaş et al. (2008) supported these results and reported that summer savory EOs reduced the development of fungal pathogens (*A. flavus*) *in vitro* even at low concentrations (25, 12.5, and 6.25 μ l/mL). The results of our study for inhibition effect are supported by several previous studies, such as Özcan and Boyraz (2000) for six moulds (*F. oxysporum* f. sp. *phaseoli*, *Macrophomina phaseoli*, *B. cinerea*, *Rhizoctonia solani*, *A. solani*, and *A. parasiticus*), Usanmaz Bozhüyük et al. (2019) for eight *Fusarium* species (*F.*

avenaceum, *F. culmorum*, *F. equiseti*, *F. graminearum*, *F. oxysporum*, *F. sambucinum*, *F. semitectum*, and *F. solani*), and Usanmaz Bozhüyük et al. (2015) for agricultural pathogenic fungus group (*Botrytis* sp., *F. equiseti*, *Nigrospora oryzae*, *P. capsici*, and *R. solani*). The fungicidal potential of essential oils has also been showed against human and plant fungal pathogens such as *A. citri*, *Alternaria* sp., *Bipolaris sorokiniana*, and *Acremonium sclerotigenum* (Yazdanpanah and Mohamadi, 2014; Mafakheri and Mirghazanfari, 2018).

ACKNOWLEDGEMENTS

In conclusion, first of all, EOs are organic and eco-friendly substances compared to antifungal pesticides. Both our study and the current literature have proven that essential oils from lavender and summer savory have an antimicrobial efficacy and can be used as an anti-*Fusarium* agent as an alternative to synthetic fungicides in the future. Therefore, in order to achieve a sustainable agriculture, these two effective essential oils could be evaluated as a potential resource in the management of main infectious fungal pathogens in plants causing damage to the agricultural ecosystem and economic loss.

REFERENCES

- Abedi-Tizaki, M., Zafari, D., Sadeghi, J. 2016. First report of *Fusarium solani* causing stem rot of *Dracaena* in Iran. *Journal of Plant Protection Research*, 56:1 210.
- Al-Naser, Z., Al-Abrass, N. 2014. Chemical composition and fungitoxic activities of *Lavandula officinalis* L. oil and comparison with synthetic fungicide on the growth some fungi in vitro. *International Journal of ChemTech Research*, 6(11): 4918-4926.
- Al-Howiriny, T.A. 2003. Composition and antimicrobial activity of essential oil of *Salvia lanigera*. *Pakistan Journal of Biological Sciences*, 6(2): 133-135.
- Al-Sadi, A.M., Al-Ghaithi, A.G., Al-Fahdi, N., Al-Yahyai, R. 2014. Characterization and pathogenicity of fungal pathogens associated with root diseases of citrus in Oman. *Int J Agric Biol*, 16: 371–376.
- Angioni, A., Barra, A., Coroneo, V., Dessi, S., Cabras, P. 2006. Chemical composition, seasonal variability, and antifungal activity of *Lavandula stoechas* L. ssp. *stoechas* essential oils from stem/leaves and flowers. *J. Agric. Food Chem*, 54(12): 4364–4370.
- Astani, A., Reichling, J., Schnitzler, P. 2011. Screening for antiviral activities of isolated compounds from essential oils. *Evidence-Based Complementary and Alternative Medicine*.
- Bammi, J., Khelifa, R., Remmal, A. 1997. Etudes de l'activité antivirale de quelques huiles essentielles. In *Proceedings of the Intern. Congr. Arom. Medicinal Plants & Essential Oils*.
- Bassolé, I.H.N., Juliani, H.R. 2012. Essential Oils in Combination and Their Antimicrobial Properties. *Molecules*. 17, 3989-4006.
- Bayaz, M. 2014. Esansiyel Yağlar: antimikrobiyal, antioksidan ve antimutajenik aktiviteleri. *Akademik Gıda*, 12(3): 45-53.
- Baytop, T. 1999. Türkiye’de Bitkilerle Tedavi. *Nobel Tıp Kitapevleri*, ISBN: 975-420-021-1 S: 332.
- Benjlali, B., Tantaoui, E.A., Ayadi, A., Ihlal, M. 1984. Method to study antimicrobial effects of essential oils: application to the antifungal activity of six Moroccan essences. *J. Food Prot.*, 47: 748–752.
- Bishop, C.D. 1995. Antiviral activity of the essential oil of *Melaleuca alternifolia* (Maiden and Betche) Cheel (tea tree) against tobacco mosaic virus. *Journal of Essential Oil Research*, 7(6): 641- 644.
- Brayford, D. 1993. The identification of *Fusarium* species. In: *Workshop Manual*. International Mycological Institute, Bakeham Lane, Egham. CAB International, Wallingford.

- Bueno, C.J., Fischer, I.H., Rosa, D.D., Firmino, A.C., Harakava, R., Oliveira, C.M.G., Furtado, E.L. 2014. *Fusarium solani* f. sp. *passiflorae*: a new forma specialis causing collar rot in yellow passion fruit. *Plant Pathol.* 63: 382–389
- Ceylan, A. 1983. Tıbbi Bitkiler-II. Ege Üniversitesi Ziraat Fakültesi Yayını No:481, Bornova-İzmir.
- Ceylan, A. 1987. Tıbbi Bitkiler 2 (Uçucu Yağ İçerenler), Ege Üniversitesi, Ziraat Fakültesi, Tarla Bitkileri Bölümü, 481:188, İzmir.
- Choi, H.J. 2018. Chemical Constituents of Essential Oils Possessing Anti-Influenza A/WS/33 Virus Activity Osong Public Health and Research Perspectives. 9(6):348–353.
- Chouhan, S., Sharma, K., Guleria, S. 2017. Antimicrobial Activity of Some Essential Oils-Present Status and Future Perspectives. *Medicines (Basel)* 4(3):58.
- Cowan, M.M. 1999. Plant products as antimicrobial agents. *Clinical Microbiology Reviews* 12(4): 564- 582.
- Çelik, E., Çelik, G.Y. 2007. Bitki uçucu yağlarının antimikrobiyal özellikleri. *Orlab On-Line Mikrobiyoloji Dergisi*, 5(2): 1-6.
- Dadalioglu, I., Evrendilek, G. 2004. Chemical compositions and antibacterial effects of essential oils of turkish oregano (*Origanum minutiflorum*), bay laurel (*Laurus nobilis*), Spanish Lavender (*Lavandula stoechas* L.), and fennel (*Foeniculum vulgare*) on Common Foodborne Pathogens. *Journal of agricultural and food chemistry*. 52. 8255.
- Deans, S.G., Sobada, K.P. 1990. Antimicrobial properties of marjoram (*Origanum marjorana* L.) volatile oil. *Flavour Fragr. J.* 5: 187-190.
- Dikbas, N., Kotan, R., Dadasoglu, F., Sahin, F. 2008. Control of *Aspergillus flavus* with essential oil and methanol extract of *Satureja hortensis*. *International Journal of Food Microbiology*, 124(2): 179-182.
- Djenane, D., Yangüela, J., Derriche, F., Bouarab, L., Roncales, P. 2012. Olive tree leaf extract; in vitro tests on staphylococcus aureus, salmonella enteritidis and pseudomonas aeruginosa; application in turkey meat. *Phytothérapie*, 10: 10–18.
- Djihane, B., Wafa, N., Elkhamssa, S., Pedro, D.H.J., Maria, E.A., Mihoub, Z.M. 2017. Chemical constituents of *Helichrysum italicum* (Roth) G. Don essential oil and their antimicrobial activity against Gram-positive and Gram-negative bacteria, filamentous fungi and *Candida albicans*, *Saudi Pharmaceutical Journal*, 25 (5): 780-787.
- El-Shazly, A., Dorai, G., Wink, M. 2002. Composition and antimicrobial activity of essential essential oils. *Int. J. Food Microbiol.* 122(1-2): 135–139.
- Farahani-Kofoet, R.D., Witzel, K., Graefe, J., Grosch, R., Zrenner, R. 2020. Species-Specific Impact of *Fusarium* Infection on the Root and Shoot Characteristics of Asparagus. *Pathogens*, 9: 509.
- Gadisa, E., Weldearegay, G., Desta, K., Tsegaye, G., Hailu, S., Jote, K., Takele, A. 2019. Combined antibacterial effect of essential oils from three most commonly used Ethiopian traditional medicinal plants on multidrug resistant bacteria. *BMC Complement Altern Med.*, 19(24).
- Gandomi, H., Misaghi, A., Basti, A.A., Bokaei, S., Khosravi, A., Abbasifar, A., Javan, A.J. 2009. Effect of *Zataria multiflora* Boiss. essential oil on growth and aflatoxin formation by *Aspergillus flavus* in culture media and cheese. *Food Chem. Toxicol.* 47(10): 2397–2400.
- Giordani, R., Regli, P., Kaloustian, J., Mikail. C., Abou, L., Portugal, H. 2004. Antifungal effect of various essential oils against *Candida albicans*. Potentiation of antifungal action of amphotericin B by essential oil from *Thymus vulgaris*. *Phytotherapy Research*, 18: 990–995.
- Grassmann, J., Elstner, E.F. 2003. Essential oils/properties and uses. *Encyclopaedia of Food Science, Food Technology and Nutrition* (Elsevier Science Ltd.), 2177-2184.

Guynot, M.E., Ramos, A.J., Seto, L., Purroy, P., Sanchis, V., Marin, S. 2003. Antifungal activity of volatile compounds generated by essential oils against fungi commonly causing deterioration of bakery products. *J. Appl. Microbiol.* 94(5): 893–899.

György, É., Laslo, É., Kuzman, I.H., András, C.D. 2020. The effect of essential oils and their combinations on bacteria from the surface of fresh vegetables. *Food Science & Nutrition*, 8: 5601–5611.

Hadian, J., Ghasemnezhad, M., Ranjbar, H., Frazane, M., Ghorbanpour, M. 2008. Antifungal potency of some essential oils in control of postharvest decay of strawberry caused by *botrytis cinerea*, *rhizopus stolonifer* and *Aspergillus niger*. *Journal of Essential Oil Bearing Plants*, 11, 553-562.

Hafizi, R., Salleh, B., Latiffah, Z. 2013. Morphological and molecular characterization of *Fusarium solani* and *F. oxysporum* associated with crown disease of oil palm. *Brazilian Journal of Microbiology*, 44(3): 959-968

Kadoglidou, K., Lagopodi, A., Karamanoli, K., Vokou, D., Bardas, G.A., Menexes, G., Constantinidou, H.I.A. 2011. Inhibitory and stimulatory effects of essential oils and individual monoterpenoids on growth and sporulation of four soil-borne fungal isolates of *Aspergillus terreus*, *Fusarium oxysporum*, *penicillium expansum*, and *Verticillium dahliae*. *European Journal of Plant Pathology*, 130: 297–309.

Kılıç, A. 2008. Uçucu Yağ Elde Etme Yöntemleri. *Bartın Orman Fakültesi Dergisi*, 10(13): 37-45.

Klaric, M.S., Kosalec, J., Mastelic, J., Pieckova, E., Pepeljak, S. 2007. Antifungal activity of thyme (*Thymus vulgaris* L.) essential oil and thymol against moulds from damp dwellings. *Lett. Appl. Microbiol.* 44(1): 36-42.

Kumar, R., Mishra, A.K., Dubey, N.K., Tripathi, Y.B. 2007. Evaluation of *Chenopodium ambrosioides* oil as a potential source of antifungal,

antiaflatoxic and antioxidant activity. *Int. J. Food Microbiol.* 115(2): 159–164.

Kurt, Ş., Uysal, A., Soylu, E.M., Kara, M., Soylu, S. 2020. Characterization and pathogenicity of *Fusarium solani* associated with dry root rot of citrus in the eastern Mediterranean region of Turkey. *Journal of General Plant Pathology*, 86: 326–332.

Lee, S.O., Choi, G.J., Jang, K.S., Lim, H.K., Cho, K.Y., Kim, J. 2007. Antifungal activity of five plant essential oils as fumigant against postharvest and soilborne plant pathogenic fungi. *Plant Pathol. Journal*, 23(2): 97-102.

Limoncu, M.E., Balcıoğlu, İ.C., Oyur, T., Zeybek, G., Zeybek, U. 2017. *In vitro* investigation of the pediculicidal activities of the volatile oil components of some medical plants raised in Turkey. *Türkiye Parazitoloj. Derg.*, 41: 208-213.

Linskens, H.F., Jackson, J.F. 1997a. *Modern Methods of Plant Analysis*, Vol. 19: *Plant Volatile Analysis*, Springer, Germany.

Lopez-Malo, A., Barreto-Valdivieso, J., Palou, E., San Martin, F. 2007. *Aspergillus flavus* growth response to cinnamon extract and sodium benzoate mixtures. *Food Control*, 18(11): 1358-1362.

Lopez-Reyes, J.G., Spadaro, D., Gullino, M.L., Garibaldi, A. 2010. Efficacy of plant essential oils on postharvest control of rot caused by fungi on four cultivars of apples in vivo. *Flavour and Fragrance Journal*, 25: 171–177.

Mafakheri, H., Mirghazanfari, S.M. 2018. Antifungal activity of the essential oils of some medicinal plants against human and plant fungal pathogens. *Cell Mol Biol.* 64(15): 13-19.

Man, A., Santacroce, L., Jacob, R., Mare, A., Man, L. 2019. Antimicrobial Activity of Six Essential Oils Against a Group of Human Pathogens: A Comparative Study. 25: 8(3).

Nguefack, J., Dongmo, J.B.L., Dakole, C.D., Leth, V., Vismer, H.F., Pedersen, J.G.T., Guemdjom, E.F.N., Mbeffo, M., Tamgue, O., Fotio, D., Zollo, P.H., Nkengfack, A.E. 2009. Food preservative potential of essential oils and fractions from *Cymbopogon citratus*, *Ocimum gratissimum* and *Thymus vulgaris* against mycotoxigenic fungi. *Int. J. Food Microbiol.*, 131(2-3): 151–156.

Nguefack, J., Dongmo, J.B.L., Dakole, C.D., Leth, V., Vismer, H.F., Pedersen, J.G.T., Guemdjom, E.F.N., Mbeffo, M., Tamgue, O., Fotio, D., Zollo, P.H., Nkengfack, A.E. 2009. Food preservative potential of essential oils and fractions from *Cymbopogon citratus*, *Ocimum gratissimum* and *Thymus vulgaris* against mycotoxigenic fungi. *Int. J. Food Microbiol.*, 131(2-3): 151–156.

Nikolis, M., Jovanovis, K.K., Markovis, T., Markovis, D., Gligorijevis, N., Radulovis, S., Sokovis, M. 2014. Chemical composition, antimicrobial, and cytotoxic properties of five Lamiaceae essential oils. *Industrial Crops and Products*, 61: 225–232.

Nostro, A., Germano, M.P., D'angelo, V., Marino, A., Cannatelli, M.A. 2000. Extraction methods and bioautography for evaluation of medicinal plant antimicrobial activity. *Lett. Appl. Microbiol.*, 30(5): 79–84.

O'Donnell, K., Ward, T.J., Robert, V.A.R.G., Crous, P.W., Geiser, D.M., Kang, S. 2015. DNA sequence-based identification of *Fusarium*: current status and future directions. *Phytoparasitica*, 43:583–595

Ondřej, M., Dostálová, R., Trojan, R. 2008. Evaluation of virulence of *Fusarium solani* isolates on pea. *Plant Protection Science*, 44(1): 9–18.

Ondřej, M., Dostálová, R., Trojan, R. 2008. Evaluation of virulence of *Fusarium solani* isolates on pea. *Plant Protect. Sci.*, 44: 9–18.

Özcan, M., Boyraz, N. 2000. Antifungal properties of some herb decoctions. *Eur Food Res Technol*, 212: 86–88.

Özgülven, M., Kırıcı, S. 1999. Farklı ekolojilerde Nane (*Mentha*) türlerinin verim ile uçucu yağ oran ve bileşenlerin araştırılması. *Tr. J. of Agriculture and Forestry*, 23: 465-472.

Özkalp, B., Özcan, M.M. 2009. Antibacterial activity of several concentrations of sater (*Satureja hortensis* L.) essential oil on spoilage and pathogenic food-related microorganisms. *World Applied Sciences Journal*, 6(4): 509-514.

Pinto, E., Salgueiro, L.R., Cavaleiro, C., Palmeira, A., Gonzalves, M.J. 2007. In vitro susceptibility of some species of yeasts and filamentous fungi to essential oils of *Salvia officinalis*. *Ind. Crop. Prod.* 26(2): 135-141.

Ramteke, P., Ghule, M., Ramteke, S. 2019. First report of *Fusarium solani* causing root rot on fenugreek (*Trigonella foenum-graecum* L.) in India. *Plant Disease*, 104.

Ramteke, P.K., Ghule, M.R., Ramteke, S.D. 2019. First report of *Fusarium solani* causing root rot on fenugreek (*Trigonella foenum-graecum* L.) in India. *Plant Disease*.

Rasooli, I., Fakoor, M.H., Yadegarinia, D., Gachkar, L., Allameh, A., Rezaei, M.B. 2008. Antimycotoxigenic characteristics of *Rosmarinus officinalis* and *Trachyspermum copticum* L. *Int. J. Food Microbiol.*, 122(1-2): 135–139.

Rasooli, I., Rezaei, M.B., Alameh, A. 2006. Growth inhibition and morphological alterations of *Aspergillus niger* by essential oils from *Thymus eriocalyx* and *Thymus x-porlock*. *Food Control*, 17(5): 359-364.

Rasooli, I., Rezaei, M.B., Alameh, A. 2006. Growth inhibition and morphological alterations of *Aspergillus niger* by essential oils from *Thymus eriocalyx* and *Thymus x-porlock*. *Food Control*, 17(5): 359-364.

Romberg, M.K., Davis, R.M. 2007. Host range and phylogeny of *Fusarium solani* f. sp. *eumartii* from potato and tomato in California. *Plant Dis.*, 91:585-592.

Rota, C., Carramiñana, J.J., Burillo, J., Herrera, A. 2004. In vitro antimicrobial activity of essential oils from aromatic plants against selected foodborne

pathogens. *Journal of Food Protection*, 67: 1252-1256.

Sahin, F., Güllüce, M., Daferera, D., Sökmen, A., Sökmen, M., Polissiou, M., Açar, G., Özer, H. 2004. Biological activities of the essential oils and methanol extract of *Origanum vulgare* ssp. *Vulgare* in the Eastern Anatolia region of Turkey. *Food Control*, 15(7): 549–557.

Saraç Sivrikaya, I., Tosun, B., Karakaya, E. 2021. *Origanum onites* L. ve *Rosmarinus officinalis* L. uçucu yağlarının kimyasal içeriklerinin ve *Fusarium solani*'ye karşı antifungal aktivitelerinin belirlenmesi. *Türk Tarım ve Doğa Bilimleri Dergisi*, 8(2): 329–335.

Saremi, H., Amiri, M.E., Ashrafi, J. 2011. Epidemiological aspects of bean decline disease caused by *Fusarium* species and evaluation of the bean resistant cultivars to disease in Northwest Iran. *African Journal of Biotechnology*, 10(66): 14954-14961.

Sartoratto, A., Machado, A.L.M., Delarmelina, C., Figueria, G.M., Duarte, M.C.T., Rehder, V.L.G. 2004. Composition and antimicrobial activity of essential oils from aromatic plants used in Brazil. *Brazilian Journal of Microbiology*, 35(4): 275-280.

Schelz, Z., Molnar, J., Hohmann, J. 2006. Antimicrobial and antiplasmid activities of essential oils. *Fitoterapia*, 77(4): 279-285.

Silva, J.K.R.D., Figueiredo, P.L.B., Byler, K.G., Setzer, W.N. 2020. Essential oils as antiviral agents. potential of essential oils to treat SARS-CoV-2 Infection: An In-Silico Investigation. *Int J Mol Sci.*, 21(10): 3426.

Śmigielski, K.B., Prusinowska, R., Krosowiak, K., Sikora, M. 2013. Comparison of qualitative and quantitative chemical composition of hydrolate and essential oils of lavender (*Lavandula angustifolia*). *J Essent Oil Res.*, 25: 291-299.

Soylu, E.M., Kurt, S., Soylu, S. 2010. In vitro and in vivo antifungal activities of the essential oils of various plants against

tomato grey mould disease agent *botrytis cinerea*. *International Journal of Food Microbiology*, 143: 183–189.

Soylu, E.M., Soylu, S., Kurt, S. 2006. Antimicrobial activities of the essential oils of various plants against tomato late blight disease agent *Phytophthora infestans*. *Mycopathologia*, 161: 119–128.

Soylu, E.M., Yiğitbaş, H., Tok, M.F., Soylu, S., Baysal, O., Kaya, A.D. 2005. Chemical Composition and Antifungal Activity of the Essential Oil of *Artemisia annua* L. Against Foliar and Soilborne Fungal Pathogens. *Z. Pflanzenk Pflanze*, 112: 229-239.

Sugiura, Y., Sugita-Konishi, Y., Kumagai, S., Reiss, E. 2003. Experimental murine hyalohyphomycosis with soil-derived isolates of *Fusarium solani*. *Med Mycol*, Jun;41(3):241-7.

Sugiura, Y., Sugita-Konishi, Y., Kumagai, S., Reis, E. 2003. Experimental murine hyalohyphomycosis with soil-derived isolates of *Fusarium solani*. *Medical Mycology*, 41: 241-247.

Şahin, F., Karaman, İ., Güllüce, M., Ögütçü, H., Şengül, M., Adıgüzel, A., Öztürk, S., Kotan, R. 2003. Evaluation of antimicrobial activities of *Satureja hortensis* L. *Journal of Ethnopharmacology*, 87(1): 61-65.

Tarek, N., Hassan, H.M., AbdelGhani, S.M.M., Radwan, I.A., Hammouda, O., El-Gendy, A.O. 2014. Comparative chemical and antimicrobial study of nine essential oils obtained from medicinal plants growing in egypt. *Beni-Suef University Journal of Basic and Applied Sciences*, 3: 149–156.

Tutun, H., Koç, N., Kart, A. 2018. Plant Essential Oils Used Against Some Bee Diseases. *Türk Tarım - Gıda Bilim ve Teknoloji dergisi*, 6(1): 34-45.

Tuxbury, K.A., Shaw, G.C., Montali, R.J., Clayton, L.A., Kwiatkowski, N.P., Dykstra, M.J., Mankowski, J.L. 2014. *Fusarium solani* species complex associated with carapace lesions and bronchitis in captive American horseshoe crabs *Limulus*

polyphemus. Dis Aquat Organ, 109:223–230.

Tzortzakis, N.K., Economakis, C.D. 2007. Antifungal activity of lemongrass (*Cymbopogon citratus* L.) essential oil against key postharvest pathogens. Innovative Food Sci. And EmergingTech., 8(2): 253–258.

Usanmaz Bozhuyuk, A., Komaki, A., Kordali, S., Ustuner, T. 2019. Assessment of the growth inhibiting effect of Satureja essential oils on different Fusarium species from Wheat. Fresenius Environmental Bulletin, 28(11): 8199-8206.

Usanmaz Bozhüyük, A., Kordali, Ş., Bölük, G. 2015. Satureja hortensis L. Uçucu Yağının Antifungal Etkisi Atatürk Univ. J. of the Agricultural Faculty, 46 (2): 107-112

Varona, S., Rodríguez-Rojo, S., Martín, Á., Cocero, M.J., Serra, A.T., Crespo, T., Duarte, C.M.M. 2013. Antimicrobial activity of lavandin essential oil formulations against three pathogenic food-borne bacteria. Industrial Crops and Products, 42: 243–250.

Vazquez, B.I., Fente, C., Franco, C.M., Vazquez, M.J., Cepeda, A. 2001. Inhibitory effects of eugenol and thymol on *Penicillium citrinum* strains in culture media and cheese. Int. J. Food Microbiol. 67(1-2): 157–163.

Viuda-Martos, M., Mohamady, M.A., Fernández-López, J., Abd ElRazik, K.A., Omer, E.A., Pérez-Alvarez, J.A., Sendra, E. 2011. In vitro antioxidant and antibacterial activities of essential oils obtained from egyptian aromatic plants. Food Control, 22: 1715–1722.

Wang, R., Wang, R., Yang, B. 2009. Extraction of essential oils from five cinnamon leaves and identification of their volatile compound compositions. Innovative Food Sci. and Emerging Tech., 10(2): 289–292.

Wendorff, W.L., Riha, W.R., Muehlenkamp, E. 1993. Growth of molds on cheese treated with heat or liquid smoke. J. Food Protect, 56(11): 963-966.

Wendorff, W.L., Wee, C. 1997. Effect of smoke and spice oils on growth of molds on oil-coated cheeses. J. Food Protect., 60(2): 153-156.

Yaseen, T., D’Onghia, A.M. 2012. Fusarium spp. associated to citrus dry root rot: an emerging issue for Mediterranean citriculture. Acta Horti, 940: 647–655

Yazdanpanah, L., Mohamadi, N. 2014. Antifungal activity of Satureja hortensis L. essential oil against *Alternaria citri*. European Journal of Experimental Biology, 4(1): 399-403

Yeşil Çelikleş, O., Hames Kocabaş, E.E., Bedir, E., Vardar Sukan, F., Özek, T., Baser, K.H.C., 2007. Antimicrobial activities of methanol extracts and essential oils of *Rosmarinus officinalis*, depending on location and seasonal variations. Food Chemistry, 100: 553-559.

Yohalem, D., Passey, T. 2011. Amendment of soils with fresh and post-extraction lavender (*Lavandula angustifolia*) and lavandin (*lavandula×intermedia*) reduce inoculum of *verticillium dahliae* and inhibit wilt in strawberry. Applied Soil Ecology, 49: 187–196.